

# Triangular lattice antiferromagnets -- open questions

*Oleg Starykh*, University of Utah

*Leon Balents*, KITP

*Hosho Katsura*, U Tokyo

*Jason Alicea*, UC Irvine

*Andrey Chubukov*, U Wisconsin

*Christian Griset*, UCSB

*Shane Head*, U Utah



Disentangling quantum many-body systems, KITP, November 11, 2010


# Outline

- Motivation ( $\text{Cs}_2\text{CuBr}_4$ ) and theoretical progress
  - ▶ crucial role of spatial anisotropy
    - $\text{Cs}_2\text{CuCl}_4$  and  $\text{Cs}_2\text{CuBr}_4$
    - organic materials ( $t, t', U\dots$ )
- Phase diagram of spatially anisotropic Heisenberg model
  - ▶ Large-S analysis of interacting spin waves near  $J'=J$
  - ▶ Approach from one dimension,  $J' \ll J$
- Open questions

# Experiment: $M=1/3$ magnetization plateau in $\text{Cs}_2\text{CuBr}_4$

★ Observed in  $\text{Cs}_2\text{CuBr}_4$  (Ono 2004, Tsuji 2007)  $J'/J = 0.5-0.75$  but not  $\text{Cs}_2\text{CuCl}_4$  [ $J'/J = 0.34$ ]

$S=1/2$



140 J. Phys. Soc. Jpn. Vol. 74 (2005) Supplement






Fig. 8. The magnetization curve and  $dM/dH$  versus  $H$  measured at  $T = 0.4$  K in magnetic fields up to 20 T




9 experimental phases

vs

3 theoretical

Fortune et al, Phys. Rev. Lett. 102, 257201 (2009)

# Progress in one dimensional $J_1$ - $J_2$ chain (zig-zag ladder)



Okunishi, Tonegawa JPSJ (2003)  
Hikihara et al PRB (2010)

Common feature: robust **plateau**;  
expect more of it in  $D=2$

**$M=1/3$  plateau**


agrees with Oshikawa, Yamanaka, Affleck  
argument (PRL 2007):

$$p S (1 - M) = \text{integer}$$

$p$  = period,  $S$  = spin,  
 $M$  = magnetization:

$$M=1/3, p=3$$

possible for all  $S$



Heirich-Meisner et al PRB (2007)

plateau is centered around  $J_2 = J_1/2$  point  
for  $S > 1/2$ ; semi-classical spin wave  
expansion is possible there (OS 2009)

# D = 2

- surprisingly complex phase diagram of spatially anisotropic triangular lattice antiferromagnet
  - no definite conclusions from numerical studies yet...
- connections with interacting boson system
  - Superfluids
  - Mott insulators
  - Supersolids

Nikuni, Shiba 1995


Heidarian, Damle 2005

Wang et al 2009

Jiang et al 2009



Heidarian, Sorella, Becca 2009

Tay, Motrunich 2010



# Spatially anisotropic model near $J' = J$

$$H = \sum_{\langle ij \rangle} J_{ij} \mathbf{S}_i \cdot \mathbf{S}_j - h \sum_i \mathbf{S}_i^z$$



Umbrella state:  
favored classically;  
energy gain  $(J-J')^2/J$

Planar states: favored by  
quantum fluctuations;  
energy gain  $J/S$

The competition is controlled by dimensionless parameter  $\delta = S(J - J')^2 / J^2$

- Technical formulation: spatial anisotropy  $J-J'$  causes softening of **interacting** (including  $1/S$  correction) spin waves


# Sketch of phase diagram ( $J-J' \ll J$ )

Questions:  
phases?  
transitions?

fully polarized state

“Exact” dilute boson calculation

Alicea, Chubukov, OS PRL 2009





*supersolid:*  $S_r^+ \sim S \cos[Q \cdot r]$  ,  $S_r^z \sim S - S \cos^2[Q \cdot r]$

# Comparison with numerics


$J_1$ - $J_2$  chain DMRG

## Variational wave function calculation

Tay, Motrunich PRB 2010




(b) "Exact" dilute boson calculation




Phase diagram for smaller  $J'/J$ ?



# $J' \ll J$ : weakly coupled Heisenberg chains in magnetic field




- non - frustrated inter-chain coupling  $\vec{S}_r \cdot \vec{S}_{r'} \rightarrow N_r^x N_{r'}^x + N_r^y N_{r'}^y + N_r^z N_{r'}^z$   
most relevant less relevant



$$2\pi R^2 < 1/(2\pi R^2)$$

spins order in the plane perpendicular to the direction of magnetic field (z):  
**umbrella / cone / spin-flop states**

- frustrated inter-chain coupling



$$\vec{S}_{x,y} \cdot (\vec{S}_{x,y+1} + \vec{S}_{x+1,y+1}) \rightarrow N_y^x \partial_x N_{y+1}^x + N_y^y \partial_x N_{y+1}^y + \sin(\delta) S_{\pi-2\delta}^z(y) S_{\pi+2\delta}^z(y+1)$$

less relevant


most relevant (small to intermediate fields)

$$1 + 2\pi R^2 > 1/(2\pi R^2)$$

★ frustration promotes collinear **SDW** order



# $J' \ll J$ limit

Katsura, OS, Balents PRB 2010




# $J' \ll J$ limit

Katsura, OS, Balents PRB 2010




# $J' \ll J$ limit to $J' = J$ point...



# Global phase diagram

*Hypothesis:* 1/3 plateau extends for all  $0 < J'/J < 1$ ;  
 other magnetization plateaux terminate above some critical  $J'/J$  ratio.



Question: how many phases are there?  
 magnetization plateaux?

# Conclusions

- ★ Magnetization plateau persists for all  $J'/J$  (?)
  - semiclassical interacting spin waves near  $J - J' \ll J$
  - 1d scaling + symmetry arguments near  $J' \ll J$
- ★ Many interesting *magnetically ordered* phases
  - global phase diagram of triangular antiferromagnet ?
  - Longitudinal SDW (?)
    - $S=1/2$  vs  $S=1$  (?)
  - plateau for ferromagnetic  $J_1$ ? [LiCuVO<sub>4</sub>]
- ★ Many open experimentally relevant questions, excellent problem for numerical studies

# Experimental relevance

