

# Fluctuations: Experiment

**Gunther Roland**  
**MIT**

*QCD in the RHIC Era*


*ITP/UCSB April 8-12 2002*

## Fluctuations: Experiment

- Survey of experimental results
  - Global fluctuations of intensive variables
  - Energy and centrality dependence
- Some mild Speculation

## Event-by-Event Fluctuations

**Are these events 'different' or 'the same'?**



- What do we mean by 'different'?
- What physics would make them more 'different' or more 'the same'?
- What does the data tell us?

ITP UCSB 4/12/02

Gunther Roland - MIT



## Global Dynamical Fluctuations



Example:  $\langle p_T \rangle_{\text{EbE}}$  Fluctuations ( $\langle p_T \rangle_{\text{EbE}} = \sum p_{Ti}$ )

- **Global:** Study variation of  $\langle p_T \rangle_{\text{EbE}}$  from event to event
- **Dynamical:** Study  $\sigma^2_{\langle p_T \rangle_{\text{EbE}}}$  relative to statistical reference

ITP UCSB 4/12/02

Gunther Roland - MIT





## Look at Event-by-Event Fluctuations in


- $K/\pi$  ratio
- $N^+, N^-$  multiplicities
- $\langle p_T \rangle$

ITP UCSB 4/12/02

Gunther Roland - MIT

## E-by-E fluctuations in the $K/\pi$ ratio

- Is strangeness enhanced in every event?
- Can we see signs of super-cooling below  $T_{\text{crit}}$ ?




ITP UCSB 4/12/02

Gunther Roland - MIT

## E-by-E fluctuations in the $K/\pi$ ratio

- Is strangeness enhanced in every event?
- Can we see signs of super-cooling below  $T_{\text{crit}}$ ?



NA49 Measurement



- Use  $dE/dx$  to identify  $\pi, K, p$  event-by-event
- Do Max Likelihood fit to extract  $K/\pi$  ratio event-by-event
- Required 2 years of detector calibration to eliminate  $dE/dx$  – multiplicity correlation

ITP UCSB 4/12/02

Gunther Roland - MIT

## E-by-E fluctuations in the $K/\pi$ ratio

NA49, PRL 86 (2001) 1965




- Dynamical fluctuations are small ( $< \sim 5\%$ )
- Compatible with resonance gas (Jeon, Koch; nuclth/9906074)
- Strangeness enhancement in every event
- Chemical freeze-out at same T in every event

ITP UCSB 4/12/02

Gunther Roland - MIT


## Charge fluctuations



- Net Charge/ $\Delta y$  Fluctuations  $\leftrightarrow$  Charge/DoF
  - Jeon, Koch hep-ph/0003168
  - Asakawa, Heinz, Mueller hep-ph/0003169
  - Change from 1-2 (QGP) to 4 (Pion Gas)
- Fluctuations frozen b/c charge conservation
  - Diffusion vs Expansion timescale
- Fluctuations of  $N^+/N^-$  ratio or  $N^+ - N^-$  difference vs statistical reference

Gunther Roland - MIT

## Charge fluctuations at RHIC: PHENIX




No sign of QGP suppression of charge fluctuations

11P UCSB 4/12/02

Gunther Roland - MIT

## More results



ITP UCSB 4/12/02

## Charge Fluctuations

- Fluctuations close to resonance gas prediction
- Little/no energy or centrality dependence
- **Where's the QGP?**
  - Diffusion wins?
  - Dilution by late stage resonance formation (Zaranek, NA49 simulations)
  - Quark coalescence (Bialas, hep-ph/0203047)

ITP UCSB 4/12/02

Gunther Roland - MIT

## $\langle p_T \rangle_{EbE}$ Fluctuations

- $p_T$  - simple observable (supposedly...)
- High statistical precision:  $\sigma_{pT, EbE} / \langle pT \rangle_{inc} < 0.1\%$
- Sensitive to many interesting scenarios
  - Critical endpoint
  - DCC production
  - Droplet formation
  - Jets
  - Any non-statistical, momentum-localized process

ITP UCSB 4/12/02

Gunther Roland - MIT

## (Too?) Many ways to measure $p_T$ Fluctuations

$$\phi_{pT} = (\langle Z \rangle / \langle M \rangle)^{1/2} - \bar{Z}^{1/2}, \text{ w/ } z = p_T - \langle p_T \rangle, Z = \sum z$$

(Gazdzicki, Mrowczynski)

$$\sigma_{dyn}^2 = (\sigma_{EbE}^2 - \sigma_{inc}^2 / \langle M \rangle) \times \langle M \rangle / (\langle M \rangle - 1)$$

(Voloshin)

$$\Delta \sigma^2 = \overline{M} \times (\langle p_T \rangle_{inc} - \langle p_T \rangle)^2 - \sigma_{EbE}^2$$

(Trainor)


Direct model comparison (NA49, WA98, PHENIX)

ITP UCSB 4/12/02

Gunther Roland - MIT

## Global $\langle p_T \rangle$ fluctuations at SPS

NA49, Phys Lett B459 (1999) 679




- Charged particles,  $4 < y_\pi < 5.5$
- $\Phi_{p_T} = 0.6 \pm 1.0 \text{ MeV}/c$
- Result consistent with statistical fluctuations only
  - Expect +5 MeV from HBT
  - Canceled by -5MeV from two-track resolution

ITP UCSB 4/12/02

Gunther Roland - MIT

## Global $\langle p_T \rangle$ fluctuations at SPS



**It's not a Gaussian....**


**...it's a Gamma-func!**

(M. Tannenbaum, Phys.Lett.B 498(2001) 29)

No. of events	98426
$\langle N \rangle$	$270.13 \pm 0.07$
$(\langle N^2 \rangle - \langle N \rangle^2)^{\frac{1}{2}}$	$23.29 \pm 0.05$
$\bar{p}_T$	$376.75 \pm 0.06 \text{ MeV}/c$
$(\bar{p}_T^2 - \bar{p}_T'^2)^{\frac{1}{2}}$	$282.2 \pm 0.1 \text{ MeV}/c$


Gunther Roland - MIT

## New Measurements



PHENIX nucl-ex/0203015

ITP UCSB 4/12/02



CERES QM'01

Gunther Roland - MIT

## Look at $\langle p_T \rangle$ Fluctuations vs

- Collision Energy
  - Critical point
- Centrality
  - Reaction mechanisms

ITP UCSB 4/12/02

Gunther Roland - MIT


## $\langle p_T \rangle_{EbE}$ Fluctuations vs $\sqrt{s}$

- Systematics not straightforward
  - Different acceptance
  - Different variables
  - Different phase-space region
- Here
  - Translate to  $\phi_{pT}$
  - Correct for acceptance assuming  $y_{corr} \gg y_{acc}$

ITP UCSB 4/12/02

Gunther Roland - MIT



## $\langle p_T \rangle_{EbE}$ Fluctuations vs $\sqrt{s}$







**This is confusing....**

ITP UCSB 4/12/02

Gunther Roland - MIT








## Learning more



- Look at p+p and p+A data
  - Connection between 'radial flow' and fluctuations
  - p+A centrality dependence
- Look at more differential measures
  - Fluctuations at low/high pT
  - 2-particle correlators

ITP UCSB 4/12/02

Gunther Roland - MIT

## Differential measures of $p_T$ Fluctuations





## Summary

- Charge and K/ $\pi$  fluctuations look like resonance gas
  - No indication for strong first order transition
  - What happens to QGP charge fluctuations?
- $\langle p_T \rangle$  fluctuations small ( $\sim 1\%$ )
  - Non-zero near mid-rapidity
  - Not suggestive of critical point
  - Possible Energy and centrality dependence
  - Are we seeing initial state scattering?
  - Alternative look at issues of 'radial flow' and 'jet quenching'

ITP UCSB 4/12/02

Gunther Roland - MIT

## CLT Variance Comparison Measures

$$\Delta \sigma_{p_t}^2 \quad (\Delta \sigma_{p_t}^2 \equiv 2\sigma_{p_t} \Delta \sigma_{p_t}) \quad \sigma_{p_t, \text{dynamical}}^2 \quad \text{'dynamical' = observed - 'statistical'}$$

difference factor  $\sim \Phi_{p_t}$

$\delta x \rightarrow a$  (single-particle 'bin'):  $\bar{m} \rightarrow \hat{m}$        $N(\Delta x)$  - total multiplicity in acceptance

bin contents:  $m(\delta x)$ ,  $n(\delta x) = N(\Delta x) / M(\Delta x, \delta x)$ ;  $\langle n(\delta x) \rangle = m(\delta x) / n(\delta x)$

$$\begin{aligned} \Delta \sigma_m^2(\delta x) &\equiv \frac{(m - n \hat{m})^2 / n}{n} - \sigma_{\hat{m}}^2 & \sigma_{m, \text{dynamical}}^2 &\equiv \overline{(\langle m \rangle - \hat{m})^2} - \sigma_{\hat{m}}^2 / \bar{N} && \text{earlier} \\ &\equiv \overline{n(\langle m \rangle - \hat{m})^2} - \sigma_{\hat{m}}^2 & \longleftrightarrow &\equiv \overline{\{ N(\langle m \rangle - \hat{m})^2 - \sigma_{\hat{m}}^2 \}} / (N-1) \\ &\equiv (N-1) \{ \overline{\langle m_i \cdot m_j \rangle_{i \neq j} - \hat{m}^2} \} & &\equiv \overline{\langle m_i \cdot m_j \rangle_{i \neq j} - \hat{m}^2} && \text{later} \\ &\equiv \{ \overline{\sum_m (\delta x)} - \overline{\sum_m (a)} \} / N(\Delta x) & & & & \\ && & & & \text{scale variation of total variance} \end{aligned}$$

These two variance comparisons *seem* algebraically similar, yet  
are the subjects of strongly conflicting statements as to performance


Trainor

4

ITP UCSB 4/12/02

Gunther Roland - MIT

## Measure Bias Observed with Data



precision ABC test - J.G. Reid

Trainor

11

ITP UCSB 4/12/02

Gunther Roland - MIT

## Charge fluctuations - D~

$$D^{\sim} = (\langle \delta R \rangle^2 \cdot \langle N_{ch} \rangle_{acc}) / (C_y \cdot C_\mu)$$

$D^{\sim} = 4$  : pion gas with global charge conservation

$D^{\sim} \approx 1$  (or 2 Heinz) : frozen QGP fluctuations with global charge conservation

$D^{\sim} \approx 2.8$  : gas with resonances and with global charge conservation

(if both particles fall in the acceptance, e.g.  $\Delta y$  window)

V. Koch, S. Jeon, hep-ph / 0003168  
 M. Bleicher, V. Koch, S. Jeon, hep-ph / 0006201

ITP UCSB 4/12/02

Gunther Roland - MIT

