

Some topics in neutrinos and nucleosynthesis


Gail McLaughlin
North Carolina State University

Collaborators: Jenni Barnes, Kelsey Lund, Evan Grohs, Jim Kneller, Matt Mumpower, Sherwood Richers, Rebecca Surman, Yonglin Zhu

Neutron star mergers: what's happening at the center?



Beautiful NSF LIGO picture



A view from perspective of micro-physics (Fig. by Perego)

Neutrino physics affects the dynamics of a BNS

Neutrino physics matters for the dynamics of the neutron star merger




Fig. From Biaottha and Rezzolla 2017

Neutrino physics changes the outcome of element synthesis

- tidal ejecta
- collisional ejecta




fig. from Bauswein et al 2013

- disk/hypermassive NS outflow
- outflow from viscous heating





fig. from Perego et al 2014

Specific examples of questions where neutrino physics is needed

Does all the r-process material in the galaxy come from neutron star mergers?


Which r-process elements do neutron star mergers make?

Electromagnetic counterpart to the neutron star merger GW signal






Material with significant opacity is the best fit to the data Slide credit: Dan Kasan Suggests lanthanides were made in the merger.

Where are the lanthanides?




The r-process, what is it?

The rapid neutron capture process of nucleosynthesis



Where are the lanthanides?



Some roles that microphysics plays


nuclear structure/reactions and the EM counterpart

- freshly synthesized nuclei decay and release energy
- some fraction of this energy thermalizes in the ejecta
- thermalized energy diffuses out at a rate determined by the opacity

two primary ways the new elements are important

- they determine the nuclear heating
- they create the opacity: more lanthanides \rightarrow higher opacity

The nuclei which decay leave an imprint on the light curve



Fission of ^{254}Cf changes the heating curve




fig. from Zhu et al 2018. The FIRE collaboration isolated the extra heating to come largely from a single nucleus.

Observable consequence






fig. from Zhu et al 2018.



How many neutrons were captured?




The weak interaction matters

How neutrinos influence nucleosynthesis

Neutrinos change the ratio of neutrons to protons





How much does it matter?



Flavor matters for nucleosynthesis

Neutrinos change the ratio of neutrons to protons



Oscillations change the spectra of ν_e s and $\bar{\nu}_e$ s

$$\nu_e \leftrightarrow \nu_\mu, \nu_\tau$$


$$\bar{\nu}_e \leftrightarrow \bar{\nu}_\mu, \bar{\nu}_\tau$$

Mergers have less ν_μ, ν_τ than ν_e and $\bar{\nu}_e$

→ oscillation reduces numbers of $\nu_e, \bar{\nu}_e$

Will neutrinos transform in mergers?

Answer, almost certainly, is yes




Neutrinos can be described by a density matrix

$$\rho = \begin{pmatrix} \rho_{ee} & \rho_{ex} \\ \rho_{ex}^* & \rho_{xx} \end{pmatrix}$$

Additional information about the phase

Tells you how likely you are to measure the neutrino as electron type

Tells you how likely you are to measure the neutrino in an x (mu or tau) state

The diagram shows a 2x2 density matrix ρ enclosed in large parentheses. The elements are ρ_{ee} , ρ_{ex} , ρ_{ex}^* , and ρ_{xx} . Three arrows point from text labels to specific elements: one from the top-right label to ρ_{ex} , one from the bottom-left label to ρ_{ee} , and one from the bottom-right label to ρ_{xx} .

Neutrinos can oscillate (flavor transform)

$$i \frac{D\rho}{Dt} = [\mathbf{H}, \rho] + i\mathbf{C}$$

$$i \frac{D\bar{\rho}}{Dt} = [\bar{\mathbf{H}}, \bar{\rho}] + i\bar{\mathbf{C}}$$

Collision
term

Convective derivative

Hamiltonian

Hamiltonian creates non-linearity

$$\mathbf{H} = \mathbf{H}_{\text{vac}} + \mathbf{H}_{\text{M}} + \mathbf{H}_{\text{SI}}$$

$$\bar{\mathbf{H}} = \mathbf{H}_{\text{vac}} - \mathbf{H}_{\text{M}} - \mathbf{H}_{\text{SI}}^*$$

$$i \frac{D\rho}{Dt} = [\mathbf{H}, \rho]$$


$$i \frac{D\bar{\rho}}{Dt} = [\bar{\mathbf{H}}, \bar{\rho}]$$

Neutrinos see a potential due to other neutrinos

Neutrinos see a potential due to the matter

Flavor and mass are not the same

Where and how these transformations might occur



$$\mathbf{H} = \mathbf{H}_{\text{vac}} + \mathbf{H}_{\text{M}} + \mathbf{H}_{\text{SI}}$$

$$\bar{\mathbf{H}} = \mathbf{H}_{\text{vac}} - \mathbf{H}_{\text{M}} - \mathbf{H}_{\text{SI}}^*$$

fig. from Malkus et al 2016

Transformation is sensitive to conditions, approximations




Fig from Deaton et al

Transformation closest to the emission: “fast flavor”

Fast flavor:


fastest transitions when inverse fluctuation wavelength (k) is similar to the difference in number density between neutrinos and antineutrinos

and

there is a “crossing”


(Sawyer, Friedland, Johns, Fuller, Balantekin, Patwardhan, Suliga and many more)

Example of a crossing



Data from Foucart, Fig from Grohs et al in prep

Crossings in BNS remnant




Ways to analyze flavor transformation

- Stability analysis → Find a growth rate
- (Toy Models)
- Particle in cell methods → track everything about every neutrino
- More approximate methods → moments

(in)stability analysis for one spatial point

15 Solar Mass, 400 ms, 465 km



Particle in cell - many neutrinos, two beam

Two beams has an analytic solution for the original growth rate (grey line)




Figure from Richers et al 2020

Toward inclusion in simulation: less exact methods: e.g. moments

What? Represent all the neutrinos at each point in space as four quantities (e.g. energy density and flux) and evolve these

Why? Possible way to eventually integrate into neutron star merger, supernova simulations

Numerical risk: Truncating an infinite tower of moments
(Fuller, Johns, Burrows, Duan ...)

Angular moments defined


$$E(t, \vec{r}, q) = \frac{1}{4\pi} \left(\frac{q}{2\pi\hbar c} \right)^3 \int d\Omega_p f(t, \vec{r}, \vec{p})$$

$$\vec{F}(t, \vec{r}, q) = \frac{1}{4\pi} \left(\frac{q}{2\pi\hbar c} \right)^3 \int d\Omega_p \hat{p} f(t, \vec{r}, \vec{p})$$

$$P(t, \vec{r}, q) = \frac{1}{4\pi} \left(\frac{q}{2\pi\hbar c} \right)^3 \int d\Omega_p \hat{p} \otimes \hat{p} f(t, \vec{r}, \vec{p})$$





Energy, flux and pressure moments

Crossings in BNS remnant




Fast flavor oscillations above a BNS merger with moments using FLASH


(Grohs et al in prep.)



Growth and saturation, BNS, moments vs PIC




Growth and saturation, BNS, moments vs PIC



Grohs et al in prep

Fourier transform BNS, moments vs PIC



Conclusions

We need to understand neutrinos in astrophysical systems to accurately predict observables including r-process

Involves solving the quantum kinetic equations in astrophysical environments

Starting to make progress on this using moment based methods

To keep mind: Astrophysical objects will make better laboratories for neutrino physics if we make progress on understanding systems with large numbers of neutrinos